

Contents

Introduction	v
1 Background	1
1.1 Raman spectroscopy	1
1.1.1 Raman effect	1
1.1.2 Scattering geometry and depolarization	5
1.1.3 Scattering cross-section	11
1.2 Supersonic jet expansion	14
1.3 Quantum chemical calculations	15
1.4 Davydov splitting	17
2 Experimental	21
2.1 <i>Curry</i> -jet setup	21
2.2 Band position error	26
2.3 Influence of the measurement conditions on the spectra	27
2.4 Chemicals and gases	30
2.5 Determination of the compound concentration	30
3 Formic acid	35
3.1 Formic acid monomer	37
3.1.1 HCOOH monomer	37
3.1.2 Monomers of the three deuterated formic acid isotopomers	49
3.1.3 Conclusions and outlook	56
3.2 Formic acid dimer	57

3.2.1	The region up to 750 cm^{-1}	62
3.2.1.1	Experimental spectra	63
3.2.1.2	Comparison with harmonic predictions	70
3.2.1.3	Anharmonicity analysis	74
3.2.1.4	Anharmonic perturbation theory	81
3.2.1.5	The lowest intramolecular modes	84
3.2.1.6	Conclusions	87
3.2.2	$750\text{-}1800\text{ cm}^{-1}$ region	88
3.2.3	Region above 1800 cm^{-1}	108
3.3	Conclusions	135
4	Acetic acid dimer	143
4.1	The region up to 200 cm^{-1}	144
4.1.1	Raman experimental results	145
4.1.2	Supramolecular stacks of dimers	148
4.1.3	Former IR studies	152
4.2	$200\text{-}1800\text{ cm}^{-1}$ region	153
4.2.1	Fundamental modes of $(\text{HAc})_2$	154
4.2.2	Fundamental modes of $(\text{DAc})_2$ and $(\text{D}_4\text{Ac})_2$	164
4.2.3	Non-fundamental modes	167
4.3	Region above 1800 cm^{-1} : $\nu(\text{O-H/D})$ and $\nu(\text{C-H/D})$ vibrations . .	171
4.4	Conclusions	183
5	Pivalic and Propiolic acid	187
5.1	Comparison of the intermolecular modes of the four acid dimers .	188
5.1.1	Effects on the band positions	188
5.1.2	Stretching force constant and dissociation energy trends .	192
5.1.3	Conclusions	197
5.2	Propiolic and pivalic acid: region above 275 cm^{-1}	198
5.2.1	Propiolic acid	198
5.2.2	Pivalic acid	200

6	Water clusters	203
6.1	Band assignment	204
6.2	Davydov splitting analysis	217
6.3	Conclusions and Outlook	218
7	Conclusions and Outlook	221
A	Bibliography	I